

**SOFT****BAIT****LUMI****FINAL<sup>®</sup>****SOFT BAIT**  
PATENT PENDING**NET WT:**  
**16 lbs (7.2 kg)****KILLS RATS & MICE***Kills Warfarin Resistant Norway Rats And House Mice*

## ACTIVE INGREDIENT:

Brodifacoum (CAS #56073-10-0): 0.0025%

OTHER INGREDIENTS\*: 99.9975%

\*Contains Denatonium Benzoate TOTAL 100.0000%

**KEEP OUT OF REACH OF CHILDREN**  
**CAUTION**

See back panels for First Aid and additional precautionary statements.

6071F

FIRST AID HAVE LABEL WITH YOU WHEN OBTAINING TREATMENT ADVICE	
If Swallowed	<ul style="list-style-type: none"><li>• Call a poison control center, doctor, or 1-877-854-2494 immediately for treatment advice.</li><li>• Have a person sip a glass of water if able to swallow.</li><li>• Do not induce vomiting unless told to do so by the poison control center or doctor.</li></ul>

NOTE TO PHYSICIAN
If swallowed, this material may reduce the clotting ability of the blood and cause bleeding. If ingested, administer Vitamin K <sub>1</sub> intramuscularly or orally. Vitamin K <sub>1</sub> is antidotal at doses of 10–20 mg (not mg/kg). Repeat as necessary based on monitoring of prothrombin times.

TREATMENT FOR PET POISONING
If animal eats bait, call veterinarian or 1-877-854-2494 at once.
NOTE TO VETERINARIAN
<b>Anticoagulant Brodifacoum:</b> For animals ingesting bait and/or showing poisoning signs (bleeding or elevated prothrombin times), give Vitamin K <sub>1</sub> at doses of 2–5 mg/kg. If needed, check prothrombin times every 3 days until values return to normal (up to 30 days). In severe cases, blood transfusions may be needed.

#### DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. **READ THIS LABEL:** Read this entire label and follow all use directions and use precautions. Use only for the sites, pests, and application methods described on this label.

**IMPORTANT:** Do not expose children, pets, or nontarget animals to rodenticides. To help to prevent exposure:

1. Store unused product out of reach of children and pets.
2. Apply bait in locations out of reach of children, pets, domestic animals and nontarget wildlife, or in tamper-resistant bait stations. These stations must be resistant to destruction by dogs and by children under six years of age, and must be used in a manner that prevents such children from reaching into bait compartments and obtaining bait. If bait can be shaken from bait stations when they are lifted, units must be secured or otherwise immobilized. Stronger bait stations are needed in areas open to hooded livestock, raccoons, bears, or other potentially destructive animals, or in areas prone to vandalism.
3. Dispose of product container and unused, spoiled, or unconsumed bait as specified on this label.

**Note:** Bait stations are mandatory for outdoor, above-ground use. Tamper-resistant bait stations must be used if children, pets, non-target mammals, or birds may access the bait.

#### USE RESTRICTIONS:

This product may only be used to control Norway rats, roof rats, and house mice *in and within* 100 feet of man-made structures constructed in a manner so as to be vulnerable to commensal rodent invasions and/or to harboring or attracting rodent infestations. Examples of such structures include homes and other permanent or temporary residences, food processing facilities, industrial and commercial buildings, trash receptacles, agricultural and public buildings, transport vehicles (ships, trains, aircraft), docks and port of terminal buildings and related structures around and associated with these sites. Fence and perimeter baiting, beyond 100 feet from a structure as defined above, is prohibited. This product must not be applied directly to food or feed crops.

This product may also be used in alleys. Do not place near or inside ventilation duct openings. Do not contaminate water, food, feedstuffs, food or feed handling equipment, or milk or meat handling equipment or surfaces that come into direct contact with food. When used in USDA inspected facilities, this product must be applied in tamper-resistant bait stations. Do not broadcast bait.

60718



#### Kills Warfarin Resistant Norway Rats And House Mice

Norway rats, roof rats, and house mice may consume a lethal dose in one night's feeding with first dead rodents appearing five or six days after feeding begins.

ACTIVE INGREDIENT:	
Brodifacoum (CAS #56073-10-0)	0.0025%
OTHER INGREDIENTS*	99.9975%
*Contains Denatonium Benzoate	TOTAL 100.0000%

## KEEP OUT OF REACH OF CHILDREN CAUTION

See side panels for First Aid and additional precautionary statements.

### STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

**Pesticide Storage:** Store only in original container in a cool, dry, place inaccessible to children and pets. Keep containers closed and away from other chemicals.

**Pesticide Disposal:** Dispose of wastes resulting from the use of this product in trash or at an approved waste disposal facility.

**Container Handling:** Nonrefillable container. Do not reuse or refill this container. [Plastic:] Offer for recycling or reconditioning; or puncture and dispose of in a sanitary landfill or by incineration. In most states, burning is not allowed. [Paper:] Dispose of empty container by placing in trash, at an approved waste disposal facility or by incineration. In most states, burning is not allowed.

**WARRANTY:** To the extent consistent with applicable law, seller makes no warranty, expressed or implied, concerning the use of this product other than indicated on the label. Buyer assumes all risk of use and/or handling of this material when such use and/or handling is contrary to label instructions.

**NET WEIGHT: 16 lbs (7.2kg)**

EPA REG. NO.: 12455-139

EPA EST. NO.: 12455-WI-1

Manufactured by:



#### DIRECTIONS FOR USE (Continued from other panel)

Do not sell this product in individual containers holding less than 16 pounds.

Burrow baiting with Final® Soft Bait is prohibited.

**SELECTION OF TREATMENT AREAS:** Determine areas where rats or mice are present and consume the bait. Generally, these areas are along walls, by gnawed burrows, in corners and concealed places, between floors and walls, or in areas where signs have been seen. Protect bait from rain and snow. Remove food as possible.

#### APPLICATION DIRECTIONS:

**RATS:** Place 6 to 28 sachets at intervals of 15 to 30 feet per placement in an uninterrupted supply of fresh bait for at least 10 days or until signs of rat activity cease.

**MICE:** Place 1 sachet per placement. Space placements at 8 to 12 foot intervals. Supply of fresh bait for at least 15 days or until signs of mouse activity cease.

**FOLLOW-UP:** Replace contaminated or spoiled bait immediately. Wear gloves when disposing of all dead, exposed animals and leftover bait. To prevent reinfestation, remove rodent food, water, and harborage as much as possible. If reinfestation occurs, repeat treatment. Where a continuous source of infestation is present, establish bait stations and replenish as needed.

#### PRECAUTIONARY STATEMENTS

**Hazards to Humans and Domestic Animals**

**CAUTION:** Harmful if swallowed. Keep away from children, domestic animals, and pets.

All handlers (including applicators) must wear shoes plus socks, and gloves when applying. Retrieve carcasses or unused bait following application of this product must be done immediately.

#### User Safety Requirements

Follow manufacturer's instruction for cleaning/maintaining PPE. If no specific instruction for cleaning/maintaining PPE is given, wash with soap and water. Washable, use detergent and hot water. Keep and wash PPE separately. Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash hands thoroughly after applying bait. Do not drink, chew gum, use tobacco or use the toilet and change into clean clothes.

#### ENVIRONMENTAL HAZARDS

This product is extremely toxic to mammals and birds. Dogs, cats and other predatory and scavenging mammals and birds might be poisoned if they feed upon animals that have eaten this bait. This pesticide is toxic to fish. Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Runoff also may be hazardous to aquatic organisms in water adjacent to treated areas. Do not contaminate water when disposing of equipment wash water or rinsate.

100512/07-12