









KILLS RATS, MICE & MEADOW VOLES*

Kills Warfarin Resistant Norway Rats
KEEP OUT OF REACH OF CHILDREN
CAUTION

See back panels for First Aid and additional precautionary statements.

ACTIVE INGREDIENT:
Bromadiologe (CAS #28772-56-7):

 Bromadiolone (CAS #28772-56-7):
 0.005%

 OTHER INGREDIENTS†:
 99.995%

 †Contains Denatonium Benzoate
 TOTAL
 100.000%

*Not permitted for use against the following species in California: Cotton rat, Eastern harvest mouse, Golden mouse, Polynesian rat, Meadow vole, White-throated woodrat, Southern plains woodrat, and Mexican woodrat.

FIRST AID

HAVE LABEL WITH YOU WHEN OBTAINING TREATMENT ADVICE

IF SWALLOWED:

- Call a poison control center, doctor, or 1-977-854-2494, or 1-900-859-7378** immediately for treatment advice.

 + Have person sip a glass of water if able to swallow.

 Do not induce wormling unless told to do so by the poison control center or doctor.

 IF ON SKIN OR CLOTHING:

- Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes.
- Call a poison control center or doctor for treatment advice.

- Hold eye open and rinse slowly and gently with water for 15–20 minutes.

 Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center, doctor, or 1-877-854-2494 immediately for treatment advice.
- ** Also call this number for information on health concerns and pesticide incidents.

NOTE TO PHYSICIAN

If swallowed or absorbed through the skin, this material may reduce the clotting ability of the blood and cause bleeding. If ingested, administer Vitamin K₁ intramuscularly or orally. Repeat as necessary based on monitoring of prothrombin times.

TREATMENT FOR PET POISONING If animal eats bait, call veterinarian at once NOTE TO VETERINARIAN

Anticoagulant Bromadiolone: For animals ingesting belt and/or showing poisoning signs (bleeding or elevated prothrombin times), give Vitamin Kr₁. If needed, check prothrombin times every 3 days until values return to normal (up to 30 days). In severe cases, blood transfusions may be needed.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. READ THIS LABEL: Read this entire label and follow all use directions and use precautions. Use only for sites, pests, and application methods described on this label.

IMPORTANT: Do not expose children, pets, or nontarget animals to rodenticides. To help to prevent exposure: Store unused product out of reach of children and pets.

- 2. A copy bat in locations out of reach of children, pels, domestic animals and nontarget wildlife, or in lamper-resistant beit stations. These stations must be resistant to destruction by dogs and by children under six years of age, and must be used in a manner that prevents such children from reaching into bait compartments and obtaining bait. If bait can be shaken from bait stations when they are lifted, units must be secured or otherwise immobilized. Stronger bait stations are needed in areas open to hoofed livestock, raccoons, bears, or other potentially destructive animals, or in

Bait stations are mandatory for outdoor, above-ground use. Tamper-resistant bait stations must be used ver children, pets, non-target mammals, or birds may have access to the balt placement location.

USE RESTRICTIONS: This product may only be used to control the following rodent pests in and around man-made OSE NEST NOT IONS: This product may only out used to drout the clowing bodies peed in all violent manness (Nut miscoulis), Norway and Flatfus nonegious), Roof and (Flatfus paths), Cotton mouse (Peromyscus gasspirus), Cotton nat; (Sigmodom hispitus), Deer mouse (Peromyscus parantizatios), Eastern harvest mouse (Flatfundordomys hurulla, Goldom mouse (Cottonomys nutual), Propession and (Flatfus evaluris), Maedow vole! (Microtius pennsylvanius), White hoted mouse (Peromyscus leuzopus), White hinterated woodrat! (Veloriona adapting), Southern plans woodrat! (Veloriona microgus), and Medician woodrat! (Veloriona adapting), Southern plans woodrat! (Veloriona microgus), and Medician woodrat! (Veloriona diputal), southern plans woodrat! (Veloriona microgus), and Medician woodrat! Neverloan micrograms to be undersible to commensal. rodent invasions and/or to harboring or attracting rodent intestations. Examples of such structures include homes and under permanent or temporary seatones, tood processing lacilities, inclusived and commercial buildings, trash ecoephales, agricultural and public buildings, transport vehicles (sitips, trains, aircraft), docks and port or terminal buildings, trains structural and public buildings, trainsport vehicles (sitips, trains, aircraft), docks and port or terminal buildings and related structures around associated with these sites. Fince and perimeter bealing, beyond 100 feet from a structure as defined above, is prohibited. This product must not be applied directly to lood or feed crops.



KILLS RATS, MICE, AND MEADOW VOLES*

Kills Warfarin Resistant Norway Rats

Norway rats, roof rats, and house mice may consume a lethal dose in one night's **feeding with** first dead rodents appearing four or five days after feeding begins.

ACTIVE INGREDIENT:

KEEP OUT OF REACH OF CHILDREN CAUTION

See side panels for First Aid and additional precautionary statements,

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by stonge or disposal.

Pesticide Stonge: Store early in original container in a cool, dry place inaccessible to children and pets. Keep containers closed and away from other chemicals.

Pesticide Disposal: Wastes resulting from the use of this product may be placed in trash or delivered to an

approved waste disposal facility.

Container Handling: Nonrefillable container. Do not reuse or refill this container. [Plastic:] Offer for recycling

or reconditioning; or puncture and dispose of in a sanitary landfill; or by incineration. In most states, burning in

WARRANTY: To the extent consistent with applicable law, seller makes no warranty, expressed or implied, erning the use of this product other than indicated on the label. Buyer assumes all risk of use and/or handling of this material when such use and/or handling is contrary to label instructions.

NET WEIGHT: 18 lbs (8.2 kg)

EPA REG. NO. 12455-79 EPA EST NO. 12455-WI-1



DIRECTIONS FOR USE (Continued from other panel)

Burrow baiting with Contrac All-Weather Blox is prohibited.

Do not place near or inside ventilation duct openings. Do not contaminate water, bood, leedstulfs, bood or leed handling equipment, or milk or meat handling equipment or surfaces that come his direct contact with bood When used in USDA inspected facilities, this product must be applied in tamper-resistant bait stations. Do not broadcast bait. Do not use this product in sewers

Do not sell this product in individual containers holding less than 16 pounds of bail

SELECTION OFTREATMENT AREAS: Determine areas where rals, mice, or meadow voles' will most likely find and consume the bat. Generally, these areas are along walls, by grawed openings, in corners and concealed places, between floors and walls, or in locations where rats, mice, or meadow votes*, or their signs have been seen. Protect bait from rain and show. Remove as much atemative food as possible.

APPLICATION DIRECTIONS:

RATS: Place 3 to 16 bat blocks (at intervals of 15 to 30 feet) per placement in infested areas. Maintain an uninterrupted supply offresh bat for at least 10 days or until signs of rat activity ocese.

MICE AND MEADOW VOLES*: Place 1 block per placement. Space placements at 8- to 12-foot intervals in infested

areas. Two blocks may be needed at points of very high activity. Maintain an uninterrupted supply of fresh bait for at least 15 days or until signs of

mouse or meadow vole 'activity cease.

FOLLOW-UP: Reptace contaminated or spoiled bait immediately.

Wearing gloves, collect and dispose of all dead, exposed animals and leftover bait. To prevent reinfestation, limit sources of rodent food, water, and harborage as much as possible. If reinfestation does occur, repeat treatment. Where a continuous source of infestation is present, establish

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION: Harmful if swallowed or absorbed through the skin. Keep away from children, domestic animals and pets. Do not get in eyes, on skin or on dothing.

All handlers (including applicators) must wear: shoes plus socks, and waterproof gloves. Any person who retrieves carcasses or unused bait following application of this product must wear gloves.



User Safety Requirements

of lower yndrumenens.

Follow manufacturer's instruction for deaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry. Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash hands thoroughly after applying bait and before eating, drinking, chewing gum, using tobacco or using the toilet and change into clean clothing

ENVIRONMENTAL HAZARDS

This product is extremely toxic to fish, birds and other wildlife. Dogs and predatory and scavenging mammals and birds might be possoned if they feed upon animals that have eaten this bail. Do not apply this product directly to water or to areas where surface water is present to it intertidal erase below the mean high water mark. Rundfl also may be hazardous to aquatic organisms in water adjacent to treated areas. Do not contaminate water when disposing of equipment wash water or rinsate.

*Not permitted for use against the following species in California: Cotton rat, Eastern harvest mouse, Golden mouse, Polynesian rat, Meadow vole, White-throated woodrat, Southern plains woodrat, and Mexican woodrat.

Product Code: CB4051 090415/09-15

587CB-9